arXiv:1202.3981v2 [math.NT] 22 Feb 2012

On Two Types of Harmonic Number Identities

M.J. Kronenburg

February 23, 2012

Abstract

Two types of finite series of products of harmonic numbers involving nonnegative integer powers are evaluated, also yielding two other important harmonic number identities. The recursion formulas for these sums are derived, which are easily translated into a computer program.

Keywords: harmonic number.
MSC 2010: 11B99

1 Definitions and Basic Identities

The generalized harmonic numbers used in this paper are:

$$H_n^{(m)} = \sum_{k=1}^{n} \frac{1}{k^m} \tag{1.1}$$

from which follows that $H_0^{(m)} = 0$. The traditional harmonic numbers are:

$$H_n = H_n^{(1)} \tag{1.2}$$

A well known identity is [2, 3, 5, 6]:

$$\sum_{k=1}^{n} \frac{1}{k} H_k = \frac{1}{2}(H_n^2 + H_n^{(2)}) \tag{1.3}$$

and [3, 4]:

$$\sum_{k=0}^{n} \frac{1}{k+1} H_k = \frac{1}{2}(H_{n+1}^2 - H_{n+1}^{(2)}) \tag{1.4}$$

Let B_n^+ be the Bernoulli number with $B_1^+ = 1/2$ instead of $B_1 = -1/2$:

$$B_n^+ = (-1)^n B_n = B_n + \delta_{n,1} \tag{1.5}$$

Then for nonnegative integer p [4]:

$$H_n^{(-p)} = \frac{1}{p+1} \sum_{k=1}^{p+1} \binom{p+1}{k} B_{p-k+1}^+ n^k \tag{1.6}$$

and for nonnegative integer p [4]:

$$\begin{aligned}&\sum_{k=0}^{n} k^p H_k^{(m)} = F(n,p,m)\\&= H_n^{(-p)} H_n^{(m)} + H_n^{(m-p)} - \frac{1}{p+1}\sum_{k=1}^{p+1}\binom{p+1}{k} B_{p-k+1}^{+} H_n^{(m-k)}\end{aligned} \tag{1.7}$$

When $a \leq b$ are two integers and $\{x_k\}$ and $\{y_k\}$ are two sequences of complex numbers, and $\{s_k\}$ the sequence of complex numbers defined by:

$$s_k = \sum_{i=a}^{k} x_i \tag{1.8}$$

then there is the following summation by parts formula [4]:

$$\sum_{k=a}^{b-1} x_k y_k = s_{b-1} y_b - \sum_{k=a}^{b-1} s_k (y_{k+1} - y_k) \tag{1.9}$$

2 First Type of Harmonic Number Identities

Theorem 2.1. *For nonnegative integer n:*

$$\sum_{k=0}^{n} H_k H_{n-k} = (n+1)[(H_{n+1}-1)^2 - H_{n+1}^{(2)} + 1] \tag{2.1}$$

Proof. With generating functions: let $[z^n]f(z)$ be the coefficient of z^n in the power series expansion of $f(z)$. Then [2]:

$$H_n = [z^n]\frac{1}{1-z}\ln(\frac{1}{1-z}) \tag{2.2}$$

The generating function of a convolution of two series is the product of their generating functions:

$$\sum_{k=0}^{n} H_k H_{n-k} = [z^n]\frac{1}{(1-z)^2}(\ln(\frac{1}{1-z}))^2 \tag{2.3}$$

These generating functions are known [2]:

$$[z^n]\frac{1}{(1-z)^2} = n+1 \tag{2.4}$$

$$[z^n](\ln(\frac{1}{1-z}))^2 = 2\begin{bmatrix} n \\ 2 \end{bmatrix}\frac{1}{n!} \tag{2.5}$$

where $\begin{bmatrix} a \\ b \end{bmatrix}$ is the Stirling number of the first kind, which in this case evaluates to [2]:

$$\begin{bmatrix} n \\ 2 \end{bmatrix} = \begin{cases} 0 & \text{if } n = 0 \\ (n-1)! H_{n-1} & \text{if } n > 0 \end{cases} \tag{2.6}$$

The product of these two generating functions results in another convolution:

$$[z^n]\frac{1}{(1-z)^2}(\ln(\frac{1}{1-z}))^2 = 2\sum_{k=1}^{n}(n-k+1)\frac{1}{k}H_{k-1} \tag{2.7}$$

which using (1.7) with $p = 0$ and $m = 1$ and (1.4) results in:

$$\begin{aligned} &2(n+1)\sum_{k=0}^{n-1}\frac{1}{k+1}H_k - 2\sum_{k=0}^{n-1}H_k \\ &= (n+1)[H_n^2 - H_n^{(2)}] - 2[nH_n - n] \end{aligned} \tag{2.8}$$

With $H_n^{(m)} = H_{n+1}^{(m)} - 1/(n+1)^m$ this results in the theorem. $\square$

Using different methods this theorem was already proved in literature [6, 7]. Now evaluating the sum directly yields:

$$\begin{aligned} \sum_{k=1}^{n}H_kH_{n-k} &= \sum_{k=1}^{n-1}\sum_{i=1}^{k}\frac{1}{i}\sum_{j=1}^{n-k}\frac{1}{j} \\ &= \sum_{i=1}^{n-1}\frac{1}{i}\sum_{k=i}^{n-1}\sum_{j=1}^{n-k}\frac{1}{j} \\ &= \sum_{i=1}^{n-1}\frac{1}{i}\sum_{j=1}^{n-i}\frac{1}{j}\sum_{k=i}^{n-j}1 \\ &= \sum_{i=1}^{n-1}\frac{1}{i}\sum_{j=1}^{n-i}\frac{1}{j}(n-i-j+1) \\ &= (n+1)\sum_{i=1}^{n-1}\frac{1}{i}H_{n-i} - \sum_{i=1}^{n-1}H_{n-i} - \sum_{i=1}^{n-1}\frac{n-i}{i} \\ &= (n+1)\sum_{i=1}^{n-1}\frac{1}{i}H_{n-i} - \sum_{i=1}^{n-1}H_i - nH_{n-1} + (n-1) \end{aligned} \tag{2.9}$$

The second sum is evaluated using (1.7) with $p = 0$ and $m = 1$, and substituting the right side of (2.8) results in the following identity:

$$\sum_{k=1}^{n}\frac{1}{k}H_{n-k} = H_n^2 - H_n^{(2)} \tag{2.10}$$

Evaluating the sum for $n+1$ and using $H_{n-k+1} = H_{n-k} + 1/(n-k+1)$ yields:

$$\begin{aligned}\sum_{k=0}^{n+1} H_k H_{n+1-k} &= \sum_{k=0}^{n} H_k H_{n-k+1} \\ &= \sum_{k=0}^{n} H_k H_{n-k} + \sum_{k=0}^{n} \frac{1}{n-k+1} H_k \\ &= \sum_{k=0}^{n} H_k H_{n-k} + \sum_{k=0}^{n} \frac{1}{k+1} H_{n-k}\end{aligned} \tag{2.11}$$

Substituting (2.1) for $n+1$ and n results in the following identity:

$$\sum_{k=0}^{n} \frac{1}{k+1} H_{n-k} = H_{n+1}^2 - H_{n+1}^{(2)} \tag{2.12}$$

Using summation by parts (1.9) with $x_k = H_{n-k}$ and $y_k = k^p H_k$, the following recursion formula results:

$$\begin{aligned}\sum_{k=0}^{n} k^p H_k H_{n-k} = \frac{1}{p+1}\{&(n+1)^p[(n+1)H_n^2 - (n-1)H_n - \frac{n}{n+1}] \\ &- (n+1)H_n \sum_{k=0}^{n}[((k+1)^p - k^p)H_k + (k+1)^{p-1}] \\ &+ \sum_{k=0}^{n}[(n-k)((k+1)^p - k^p) + pk^p]H_k H_{n-k} \\ &+ \sum_{k=0}^{n} k[(k+1)^p - k^p + (n-k+1)^{p-1}]H_k \\ &+ \sum_{k=0}^{n} k(k+1)^{p-1}\}\end{aligned} \tag{2.13}$$

For $p = 0$ this formula yields (2.1) using (2.12). Letting $R(n,0)$ be (2.1) and using $F(n,p,m)$ from (1.7) evaluation of this formula results in the following recursion formula for integer $p > 0$:

$$\begin{aligned}\sum_{k=0}^{n} k^p H_k H_{n-k} &= R(n,p) \\ &= \frac{1}{p+1}\{-n(n+1)^p H_{n+1} \\ &+ \sum_{k=0}^{p-1} \binom{p}{k}[(n - \frac{k}{p-k+1})R(n,k) \\ &+ (1 + (-1)^k \frac{p-k}{p}(n+1)^{p-k-1})F(n,k+1,1) \\ &+ \frac{p-k}{p} H_n^{(-k-1)}]\}\end{aligned} \tag{2.14}$$

For a list of the resulting expressions up to $p = 5$ see section (4) below.

3 Second Type of Harmonic Number Identities

Theorem 3.1. *For nonnegative integer n:*

$$\sum_{k=0}^{n} H_k^2 = (n+1)H_{n+1}^2 - (2n+3)H_{n+1} + 2(n+1) \tag{3.1}$$

Proof.

$$\begin{aligned}
\sum_{k=0}^{n} H_k^2 &= \sum_{k=1}^{n}\sum_{i=1}^{k} \frac{1}{i} H_k \\
&= \sum_{i=1}^{n} \frac{1}{i} \sum_{k=i}^{n} H_k \\
&= \sum_{i=1}^{n} \frac{1}{i} (\sum_{k=1}^{n} H_k - \sum_{k=1}^{i-1} H_k) \\
&= H_n[(n+1)H_{n+1} - (n+1)] - \sum_{i=1}^{n} \frac{1}{i}(iH_i - i) \\
&= H_n[(n+1)H_{n+1} - (n+1)] - (n+1)H_{n+1} + 2n + 1 \\
&= (n+1)H_{n+1}^2 - (2n+3)H_{n+1} + 2(n+1)
\end{aligned} \tag{3.2}$$

□

Using summation by parts (1.9) with $x_k = H_k$ and $y_k = k^p H_k$, the following recursion formula results:

$$\begin{aligned}
\sum_{k=0}^{n} k^p H_k^2 = \frac{1}{p+1}\{&(n+1)^p[(n+1)H_n^2 - (n-1)H_n - \frac{n}{n+1}] \\
&- \sum_{k=0}^{n} [(k+1)((k+1)^p - k^p) - pk^p] H_k^2 \\
&- \sum_{k=0}^{n} [(k+1)^p - k((k+1)^p - k^p)] H_k \\
&+ \sum_{k=0}^{n} k(k+1)^{p-1}\}
\end{aligned} \tag{3.3}$$

For $p = 0$ this formula yields (3.1). Letting $S(n, 0)$ be (3.1) and using $F(n, p, m)$ from (1.7) evaluation of this formula results in the following recursion formula for integer

$p > 0$:

$$\begin{aligned}\sum_{k=0}^{n} k^p H_k^2 &= S(n,p)\\ &= \frac{1}{p+1}\{(n+1)[(n+1)^p(H_{n+1}^2 - H_{n+1}) - H_{n+1} + 1]\\ &- \sum_{k=0}^{p-1}\binom{p}{k}[(1+\frac{k}{p-k+1})S(n,k)\\ &+(\frac{p-k}{k+1}-1)F(n,k+1,1)\\ &-\frac{p-k}{p}H_n^{(-k-1)}]\}\end{aligned} \tag{3.4}$$

For a list of the resulting expressions up to $p = 5$ see section (4) below.
The following harmonic number identities can be expressed as a sum of this type:

$$\begin{aligned}\sum_{k=0}^{n} k^p H_{n-k}^2 &= \sum_{k=0}^{n}(n-k)^p H_k^2\\ &= \sum_{k=0}^{p}(-1)^k\binom{p}{k}n^{p-k}S(n,k)\end{aligned} \tag{3.5}$$

For a list of the resulting expressions up to $p = 5$ see section (4) below.

4 Examples

$$\sum_{k=1}^{n}\frac{1}{k}H_k = \frac{1}{2}(H_n^2 + H_n^{(2)}) \tag{4.1}$$

$$\sum_{k=0}^{n}\frac{1}{k+1}H_k = \frac{1}{2}(H_{n+1}^2 - H_{n+1}^{(2)}) \tag{4.2}$$

$$\sum_{k=1}^{n}\frac{1}{k}H_{n-k} = H_n^2 - H_n^{(2)} \tag{4.3}$$

$$\sum_{k=0}^{n}\frac{1}{k+1}H_{n-k} = H_{n+1}^2 - H_{n+1}^{(2)} \tag{4.4}$$

Formula (1.6):

$$H_n^{(0)} = n \tag{4.5}$$

$$H_n^{(-1)} = \frac{1}{2}n(n+1) \tag{4.6}$$

$$H_n^{(-2)} = \frac{1}{6}n(n+1)(2n+1) \tag{4.7}$$

$$H_n^{(-3)} = \frac{1}{4}n^2(n+1)^2 \tag{4.8}$$

$$H_n^{(-4)} = \frac{1}{30}n(n+1)(2n+1)(3n^2+3n-1) \tag{4.9}$$

$$H_n^{(-5)} = \frac{1}{12}n^2(n+1)^2(2n^2+2n-1) \tag{4.10}$$

Formula (2.14):

$$\sum_{k=0}^{n} H_k H_{n-k} = (n+1)[(H_{n+1}-1)^2 - H_{n+1}^{(2)} + 1] \tag{4.11}$$

$$\sum_{k=0}^{n} kH_k H_{n-k} = H_n^{(-1)}[(H_{n+1}-1)^2 - H_{n+1}^{(2)} + 1] \tag{4.12}$$

$$\begin{aligned}\sum_{k=0}^{n} k^2 H_k H_{n-k} = H_n^{(-2)}[H_{n+1}^2 - H_{n+1}^{(2)}] &- \frac{1}{18}n(n+1)(13n+5)H_{n+1} \\ &+ \frac{1}{108}n(n+1)(71n+37)\end{aligned} \tag{4.13}$$

$$\begin{aligned}\sum_{k=0}^{n} k^3 H_k H_{n-k} = H_n^{(-3)}[H_{n+1}^2 - H_{n+1}^{(2)}] &- \frac{1}{12}n^2(n+1)(7n+5)H_{n+1} \\ &+ \frac{1}{72}n^2(n+1)(35n+37)\end{aligned} \tag{4.14}$$

$$\begin{aligned}\sum_{k=0}^{n} k^4 H_k H_{n-k} &= H_n^{(-4)}[H_{n+1}^2 - H_{n+1}^{(2)}] \\ &\quad - \frac{1}{900}n(n+1)(447n^3+468n^2+17n-32)H_{n+1} \\ &\quad + \frac{1}{54000}n(n+1)(20739n^3+33066n^2+4129n-3934)\end{aligned} \tag{4.15}$$

$$\begin{aligned}\sum_{k=0}^{n} k^5 H_k H_{n-k} &= H_n^{(-5)}[H_{n+1}^2 - H_{n+1}^{(2)}] \\ &\quad - \frac{1}{360}n^2(n+1)(157n^3+218n^2+17n-32)H_{n+1} \\ &\quad + \frac{1}{21600}n^2(n+1)(6839n^3+14566n^2+4129n-3934)\end{aligned} \tag{4.16}$$

Formula (3.4):

$$\sum_{k=0}^{n} H_k^2 = (n+1)H_{n+1}^2 - (2n+3)H_{n+1} + 2(n+1) \tag{4.17}$$

$$\sum_{k=0}^{n} kH_k^2 = H_n^{(-1)}H_{n+1}^2 - \frac{1}{2}(n^2+n-1)H_{n+1} + \frac{1}{4}(n+1)(n-2) \tag{4.18}$$

$$\sum_{k=0}^{n} k^2 H_k^2 = H_n^{(-2)} H_{n+1}^2 - \frac{1}{18}(4n^3+9n^2+5n+3)H_{n+1} + \frac{1}{108}(n+1)(8n^2+n+18) \tag{4.19}$$

$$\begin{aligned}\sum_{k=0}^{n} k^3 H_k^2 = H_n^{(-3)} H_{n+1}^2 &- \frac{1}{24} n(n+1)(n+2)(3n+1)H_{n+1} \\ &+ \frac{1}{288} n(n+1)(9n^2+13n+14)\end{aligned} \tag{4.20}$$

$$\begin{aligned}\sum_{k=0}^{n} k^4 H_k^2 &= H_n^{(-4)} H_{n+1}^2 \\ &- \frac{1}{900}(72n^5+315n^4+410n^3+135n^2-32n-30)H_{n+1} \\ &+ \frac{1}{54000}(n+1)(864n^4+2241n^3+2629n^2+1466n-1800)\end{aligned} \tag{4.21}$$

$$\begin{aligned}\sum_{k=0}^{n} k^5 H_k^2 &= H_n^{(-5)} H_{n+1}^2 \\ &- \frac{1}{360} n(n+1)(n+2)(2n+1)(10n^2+19n-9)H_{n+1} \\ &+ \frac{1}{21600} n(n+1)(200n^4+736n^3+1159n^2+971n-366)\end{aligned} \tag{4.22}$$

Formula (3.5):

$$\sum_{k=0}^{n} H_{n-k}^2 = (n+1)H_{n+1}^2 - (2n+3)H_{n+1} + 2(n+1) \tag{4.23}$$

$$\sum_{k=0}^{n} k H_{n-k}^2 = H_n^{(-1)} H_{n+1}^2 - \frac{1}{2}(3n^2+5n+1)H_{n+1} + \frac{1}{4}(n+1)(7n+2) \tag{4.24}$$

$$\begin{aligned}\sum_{k=0}^{n} k^2 H_{n-k}^2 = H_n^{(-2)} H_{n+1}^2 &- \frac{1}{18}(2n+1)(11n^2+17n+3)H_{n+1} \\ &+ \frac{1}{108}(n+1)(170n^2+109n+18)\end{aligned} \tag{4.25}$$

$$\begin{aligned}\sum_{k=0}^{n} k^3 H_{n-k}^2 = H_n^{(-3)} H_{n+1}^2 &- \frac{1}{24} n(n+1)(25n^2+37n+10)H_{n+1} \\ &+ \frac{1}{288} n(n+1)(415n^2+427n+130)\end{aligned} \tag{4.26}$$

$$\begin{aligned}&\sum_{k=0}^{n} k^4 H_{n-k}^2 = H_n^{(-4)} H_{n+1}^2 \\ &- \frac{1}{900}(2n+1)(411n^4+1002n^3+679n^2+28n-30)H_{n+1} \\ &+ \frac{1}{54000}(n+1)(72114n^4+103491n^3+46129n^2+1466n-1800)\end{aligned} \tag{4.27}$$

$$
\begin{aligned}
&\sum_{k=0}^{n} k^5 H_{n-k}^2 = H_n^{(-5)} H_{n+1}^2 \\
&- \frac{1}{120} n(n+1)(98n^4 + 236n^3 + 159n^2 + n - 14) H_{n+1} \\
&+ \frac{1}{21600} n(n+1)(26978n^4 + 49996n^3 + 29599n^2 + 1961n - 3234)
\end{aligned}
\tag{4.28}
$$

5 Computer Program

The Mathematica® [8] program used to compute the expressions is given below, which should be added to the program listed in [4].

```
HarmSumR[0]=(n+1)((HarmonicNumber[n+1]-1)^2-HarmonicNumber[n+1,2]+1);
HarmSumR[p_]:=HarmSumR[p]=Simplify[1/(p+1)
 (-n(n+1)^p HarmonicNumber[n+1]
 +Sum[Binomial[p,k]((n-k/(p-k+1))HarmSumR[k]
 +(1+(-1)^k(p-k)/p(n+1)^(p-k-1))HarmSumF[k+1,1]
 +(p-k)/p HarmFun[-k-1]),{k,0,p-1}])]
HarmonicSumR[p_]:=Module[{t=HarmTable[2],u},
 u=Factor[CoefficientArrays[HarmSumR[p],t]];
 u[[1]]+Dot[u[[2]],t]+Dot[Dot[u[[3]],t],t]]
HarmSumS[0]=(n+1)HarmonicNumber[n+1]^2-(2n+3)HarmonicNumber[n+1]
 +2(n+1);
HarmSumS[p_]:=HarmSumS[p]=Simplify[1/(p+1)
 ((n+1)((n+1)^p(HarmonicNumber[n+1]^2-HarmonicNumber[n+1])
  -HarmonicNumber[n+1]+1)
 -Sum[Binomial[p,k]((1+k/(p-k+1))HarmSumS[k]
 +((p-k)/(k+1)-1)HarmSumF[k+1,1]
 -(p-k)/p HarmFun[-k-1]),{k,0,p-1}])]
HarmonicSumS[p_]:=Module[{t=HarmTable[1],u},
 u=Factor[CoefficientArrays[HarmSumS[p],t]];
 u[[1]]+Dot[u[[2]],t]+Dot[Dot[u[[3]],t],t]]
HarmSumT[p_]:=Simplify[
 Sum[(-1)^k Binomial[p,k]n^(p-k)HarmSumS[k],{k,0,p}]]
HarmonicSumT[p_]:=Module[{t=HarmTable[1],u},
 u=Factor[CoefficientArrays[HarmSumT[p],t]];
 u[[1]]+Dot[u[[2]],t]+Dot[Dot[u[[3]],t],t]]

(* Compute some examples *)
HarmonicSumR[3]//TraditionalForm
HarmonicSumS[4]//TraditionalForm
HarmonicSumT[2]//TraditionalForm
```

References

[1] M. Abramowitz, I.A. Stegun, *Handbook of Mathematical Functions*, Dover, 1970.

[2] R.L. Graham, D.E. Knuth, O. Patashnik, *Concrete Mathematics, A Foundation for Computer Science*, 2nd ed., Addison-Wesley, 1994.

[3] D.E. Knuth, *The Art of Computer Programming, Volume 1: Fundamental Algorithms*, 3rd ed., Addison-Wesley, 1997.

[4] M.J. Kronenburg, Some Generalized Harmonic Number Identities, `arXiv:1103.5430 [math.NT]`

[5] D.Y. Savio, E.A. Lamagna, S.M. Liu, Summation of Harmonic Numbers, in *Computers and Mathematics*, eds. E. Kaltofen, S.M. Watt, Springer, 1989.

[6] J. Spieß, Some Identities Involving Harmonic Numbers, *Math. Comp.* 55 (1990) 839-863.

[7] C. Wei, D. Gong, Q. Wang, Chu-Vandermonde Convolution and Harmonic Number Identities, `arXiv:1201.0420 [math.CO]`

[8] S. Wolfram, *The Mathematica Book*, 5th ed., Wolfram Media, 2003.